

Summary of Antimicrobial Prescribing Guidance – Managing Common Infections

For all PHE guidance, follow [PHE's principles of treatment](#). It is important to use antimicrobials prudently to support the national 5 year plan on [Tackling Antimicrobial Resistance](#) (click for details)

Aims

- To provide a simple, effective, economical and empirical approach to the treatment of common infections.
- To minimise the emergence of bacterial resistance in the community.

Principles of Treatment

- This guidance is based on the best available evidence, however professional judgement should be used and patients should be involved in the decision.
- Prescribe an antibiotic only when there is likely to be a clear clinical benefit.
- It is important to initiate antibiotics as soon as possible in severe infection.
- Where an empirical therapy has failed or special circumstances exist, microbiological advice can be obtained via the BVH switchboard 01253 300000
- Consider a 'No' or 'Back-up/Delayed', antibiotic strategy for acute self-limiting upper respiratory tract infections and mild UTI symptoms.
- Limit prescribing over the telephone to exceptional cases.
- Use simple generic antibiotics if possible. Avoid broad spectrum antibiotics (eg. co-amoxiclav, quinolones and cephalosporins) when narrow spectrum antibiotics remain effective, as they increase the risk of Clostridium difficile, MRSA and resistant UTIs.
- A dose and duration of treatment for adults is usually suggested, but may need modification for age, weight and renal function. In severe or recurrent cases consider a higher dose or longer course.
- Lower threshold for antibiotics in immunocompromised or those with multiple morbidities; culture and seek advice.
- Avoid widespread use of topical antibiotics (especially those agents also available as systemic preparations, e.g. fusidic acid).

Summary of antimicrobial prescribing guidance – managing common infections (October 2020)

Review October 2021



- In pregnancy take specimens to inform treatment, use this guidance alternative or seek expert advice. Penicillins, cephalosporins and erythromycin are not associated with increased risks. If possible avoid tetracyclines, aminoglycosides, quinolones, azithromycin, clarithromycin, high dose metronidazole (2 g stat) unless benefit outweighs risks. Short-term use of nitrofurantoin is not expected to cause foetal problems (theoretical risk of neonatal haemolysis). Trimethoprim is also unlikely to cause problems unless poor dietary folate intake or taking another folate antagonist e.g. antiepileptics.
- This guidance should not be used in isolation; it should be supported with patient information about back-up/delayed antibiotics, infection severity and usual duration, clinical staff education, and audits. Materials are available on the RCGP TARGET website.
- See BNF for appropriate use and dosing in specific populations, for example, hepatic impairment, renal impairment, pregnancy and breastfeeding.

Key:  Click to access doses for children  Click to access NICE's printable visual summary

*Safety advice on quinolones - consider MHRA/CHM advice when prescribing fluoroquinolones: small increased risk of aortic aneurysm and dissection and tendon damage







Jump to section on:

[Upper RTI](#)
[Lower RTI](#)
[UTI](#)
[Meningitis](#)
[GI](#)
[Genital](#)
[Skin](#)
[Eye](#)
[Dental](#)

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
▼ Upper respiratory tract infections						
Acute sore throat NICE Public Health England Last updated: Jan 2018	Advise paracetamol, or if preferred and suitable, ibuprofen for pain. Medicated lozenges may help pain in adults. Use FeverPAIN or Centor to assess symptoms: FeverPAIN 0-1 or Centor 0-2: no antibiotic; FeverPAIN 2-3: no or back-up antibiotic; FeverPAIN 4-5 or Centor 3-4: immediate or back-up antibiotic. Systemically very unwell or high risk of complications: immediate antibiotic. *5 days of phenoxymethylpenicillin may be enough for symptomatic cure; but a 10-day course may increase the chance of microbiological cure. <i>For detailed information click the visual summary icon.</i>	First choice: phenoxymethylpenicillin	500mg QDS or 1000mg BD		5 to 10 days*	
		Penicillin allergy: clarithromycin OR	250mg to 500mg BD		5 days	
		erythromycin (preferred if pregnant)	250mg to 500mg QDS or 500mg to 1000mg BD		5 days	




Summary of antimicrobial prescribing guidance – managing common infections (October 2020)


Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Influenza Public Health England Last updated: Feb 2019	Annual vaccination is essential for all those 'at risk' of influenza. ^{1D} Antivirals are not recommended for healthy adults. ^{1D,2A+} Treat 'at risk' patients with 5 days oseltamivir 75mg BD, ^{1D} when influenza is circulating in the community, and ideally within 48 hours of onset (36 hours for zanamivir treatment in children), ^{1D,3D} or in a care home where influenza is likely. ^{1D,2A+} At risk: pregnant (and up to 2 weeks post-partum); children under 6 months; adults 65 years or older; chronic respiratory disease (including COPD and asthma); significant cardiovascular disease (not hypertension); severe immunosuppression; chronic neurological, renal or liver disease; diabetes mellitus; morbid obesity (BMI>40). ^{4D} See the PHE Influenza guidance for the treatment of patients under 13 years. ^{4D} In severe immunosuppression, or oseltamivir resistance, use zanamivir 10mg BD ^{5A+,6A+} (2 inhalations twice daily by diskhaler for up to 10 days) and seek advice. ^{4D} Access supporting evidence and rationales on the PHE website .					
Scarlet fever (GAS) Public Health England Last updated: Oct 2018	Prompt treatment with appropriate antibiotics significantly reduces the risk of complications. ^{1D} Vulnerable individuals (immunocompromised, the comorbid, or those with skin disease) are at increased risk of developing complications. ^{1D}	Phenoxyethylpenicillin ^{2D}	500mg QDS ^{2D}		10 days ^{3A+,4A+,5A+}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		Penicillin allergy: clarithromycin ^{2D}	250mg to 500mg BD ^{2D}		5 days ^{2D,5A+}	
		Optimise analgesia ^{2D} and give safety netting advice				
Acute otitis media NICE Public Health England Last updated: Feb 2018	Regular paracetamol or ibuprofen for pain (right dose for age or weight at the right time and maximum doses for severe pain). Otorrhoea or under 2 years with infection in both ears: no, back-up or immediate antibiotic. Otherwise: no or back-up antibiotic. Systemically very unwell or high risk of complications: immediate antibiotic. <i>For detailed information click on the visual summary.</i>	First choice: amoxicillin	-		5 to 7 days	
		Penicillin allergy: clarithromycin OR erythromycin (preferred if pregnant)	-		5 to 7 days	
		Second choice: co-amoxiclav	-		5 to 7 days	
Acute otitis externa Public Health England Last updated: Nov 2017	First line: analgesia for pain relief, ^{1D,2D} and apply localised heat (such as a warm flannel). ^{2D} Second line: topical acetic acid or topical antibiotic +/- steroid: similar cure at 7 days. ^{2D,3A+,4B-} If cellulitis or disease extends outside ear canal, or systemic signs of infection, start oral flucloxacillin and refer to exclude malignant otitis externa. ^{1D}	Second line: topical acetic acid 2% ^{2D,4B-} OR topical neomycin sulphate with corticosteroid ^{2D,5A-} (consider safety issues if perforated tympanic membrane) ^{6B-}	1 spray TDS ^{5A-}		7 days ^{5A}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
			3 drops TDS ^{5A-}		7 days (min) to 14 days (max) ^{3A+}	
		If cellulitis: flucloxacillin ^{7B+}	250mg QDS ^{2D}		7 days ^{2D}	


Summary of antimicrobial prescribing guidance – managing common infections (October 2020)



Review October 2021


Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
			If severe: 500mg QDS ^{2D}			
Sinusitis NICE Public Health England Last updated: Oct 2017	Advise paracetamol or ibuprofen for pain. Little evidence that nasal saline or nasal decongestants help, but people may want to try them. Symptoms for 10 days or less: no antibiotic. Symptoms with no improvement for more than 10 days: no antibiotic or back-up antibiotic depending on likelihood of bacterial cause. Consider high-dose nasal corticosteroid (if over 12 years). Systemically very unwell or high risk of complications: immediate antibiotic. <i>For detailed information click on the visual summary.</i>	First choice: phenoxymethylpenicillin	500mg QDS		5 days	
		Penicillin allergy: doxycycline (not in under 12s) OR clarithromycin OR	200mg on day 1, then 100mg OD		5 days	
		erythromycin (preferred if pregnant)	500mg BD or 250 to 500mg QDS or 500 to 1000mg BD			
		Second choice or first choice if systemically very unwell or high risk of complications: co-amoxiclav	500/125mg TDS		5 days	
▼ Lower respiratory tract infections						
*Note: Low doses of penicillins are more likely to select for resistance. ^{1D} Do not use fluoroquinolones (ciprofloxacin, ofloxacin) first line because they may have long-term side effects and there is poor pneumococcal activity. ^{2B--3D-} Reserve all fluoroquinolones (including levofloxacin) for proven resistant organisms. ^{1D}						
Acute exacerbation of COPD NICE	Many exacerbations are not caused by bacterial infections so will not respond to antibiotics. Consider an antibiotic, but only after taking into account severity of symptoms (particularly sputum colour changes and increases in volume or thickness), need for hospitalisation, previous exacerbations, hospitalisations and risk of complications, previous sputum culture and susceptibility results, and risk of resistance with	First choice: amoxicillin OR	500mg TDS (see BNF for severe infection)	-	5 days	
		doxycycline OR	200mg on day 1, then 100mg OD (see BNF for severe infection)	-		
		clarithromycin	500mg BD	-		


Infection	Key points	Medicine	Doses		Length	Visual summary	
			Adult	Child			
Public Health England Last updated: Dec 2018	repeated courses. Some people at risk of exacerbations may have antibiotics to keep at home as part of their exacerbation action plan. <i>For detailed information click on the visual summary. See also the NICE guideline on COPD in over 16s.</i> <i>Please note - NICE recommendations adapted on local decisions</i>	Second choice: use alternative first choice					
		Alternative choice (if person at higher risk of treatment failure): co-amoxiclav OR	500/125mg TDS	-	5 days		
		Non-serious penicillin allergy: Cefixime	200mg BD	-			
		Severe penicillin allergy: co-trimoxazole (consider safety issues)	960mg BD				
		levofloxacin (with specialist advice) if co-amoxiclav, cefixime or co-trimoxazole cannot be used; consider safety issues)	Refer to microbiologist	-			
		IV antibiotics- Refer to/Contact Community IV Team – 01253 951223					
Acute exacerbation of bronchiectasis (non-cystic fibrosis)	Send a sputum sample for culture and susceptibility testing. Offer an antibiotic. When choosing an antibiotic, take account of severity of symptoms and risk of treatment failure. People who may be at higher risk of treatment failure include people who've had	First choice empirical treatment: amoxicillin (preferred if pregnant) OR	500mg TDS		7 to 14 days		
		doxycycline (not in under 12s) OR	200mg on day 1, then 100mg OD				
		clarithromycin	500mg BD				

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
<p>NICE</p> <p>Public Health England</p> <p>Last updated: Dec 2018</p>	<p>repeated courses of antibiotics, a previous sputum culture with resistant or atypical bacteria, or a higher risk of developing complications.</p> <p>Course length is based on severity of bronchiectasis, exacerbation history, severity of exacerbation symptoms, previous culture and susceptibility results, and response to treatment.</p> <p>Do not routinely offer antibiotic prophylaxis to prevent exacerbations.</p> <p>Seek specialist advice for preventing exacerbations in people with repeated acute exacerbations. This may include a trial of antibiotic prophylaxis after a discussion of the possible benefits and harms, and the need for regular review.</p> <p><i>For detailed information click on the visual summary.</i></p>	<p>Alternative choice (if person at higher risk of treatment failure) empirical treatment: co-amoxiclav OR</p>	500/125mg TDS		7 to 14 days	
		<p>Levofloxacin* (adults only: with specialist advice if co-amoxiclav cannot be used; consider safety issues) OR</p>	Refer to microbiologist			
		<p>ciprofloxacin (children only: with specialist advice if co-amoxiclav cannot be used; consider safety issues)</p>	-			
		<p>IV antibiotics – Refer to/Contact Community IV Team – 01253 951223</p>				
<p>When current susceptibility data available: choose antibiotics accordingly</p>						

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Acute cough NICE Public Health England Last updated: Feb 2019	Some people may wish to try honey (in over 1s), the herbal medicine pelargonium (in over 12s), cough medicines containing the expectorant guaifenesin (in over 12s) or cough medicines containing cough suppressants, except codeine, (in over 12s). These self-care treatments have limited evidence for the relief of cough symptoms. Acute cough with upper respiratory tract infection: no antibiotic. Acute bronchitis: no routine antibiotic. Acute cough and higher risk of complications (at face-to-face examination): immediate or back-up antibiotic. Acute cough and systemically very unwell (at face to face examination): immediate antibiotic. Higher risk of complications includes people with pre-existing comorbidity; young children born	Adults first choice: doxycycline	200mg on day 1, then 100mg OD	-	5 days	
		Adults alternative first choices: amoxicillin (preferred if pregnant) OR clarithromycin OR	500mg TDS	-		
		erythromycin (preferred if pregnant)	250mg to 500mg QDS or 500mg to 1000mg BD	-		
		Children first choice: amoxicillin	-		5 days	
		Children alternative first choices: clarithromycin OR erythromycin OR	-			
			-			


Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
	<p>prematurely; people over 65 with 2 or more of, or over 80 with 1 or more of: hospitalisation in previous year, type 1 or 2 diabetes, history of congestive heart failure, current use of oral corticosteroids.</p> <p>Do not offer a mucolytic, an oral or inhaled bronchodilator, or an oral or inhaled corticosteroid unless otherwise indicated.</p> <p><i>For detailed information click on the visual summary. See also the NICE guideline on pneumonia for prescribing antibiotics in adults with acute bronchitis who have had a C-reactive protein (CRP) test (CRP<20mg/l: no routine antibiotic, CRP 20 to 100mg/l: back-up antibiotic, CRP>100mg/l: immediate antibiotic).</i></p>	doxycycline (not in under 12s)	-			
<p>Community-acquired pneumonia</p> <p>NICE</p> <p>Public Health England</p>	<p>Assess severity in adults based on clinical judgement guided by mortality risk score (CRB65 or CURB65). See the NICE guideline on pneumonia for full details:</p> <p>low severity – CRB65 0 or CURB65 0 or 1 moderate severity – CRB65 1 or 2 or CURB65 2 High severity – CRB65 3 or 4 or CURB65 3 to 5. 1 point for each parameter: confusion, (urea >7 mmol/l), respiratory rate ≥30/min, low systolic (<90 mm Hg) or diastolic (≤60 mm Hg) blood</p>	<p>First choice (low severity in adults or non-severe in children): amoxicillin</p> <p>Alternative first choice (low severity in adults or non-severe in children): doxycycline (not in under 12s) OR clarithromycin OR erythromycin (in pregnancy)</p>	<p>500mg TDS (higher doses can be used, see BNF)</p> <p>200mg on day 1, then 100mg OD</p> <p>500mg BD</p> <p>500mg QDS</p>		5 days*	

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Last updated: Sept 2019	<p>pressure, age ≥65.</p> <p>Assess severity in children based on clinical judgement.</p> <p>Offer an antibiotic. Start treatment as soon as possible after diagnosis, within 4 hours (within 1 hour if sepsis suspected and person meets any high risk criteria – see the NICE guideline on sepsis).</p> <p>When choosing an antibiotic, take account of severity, risk of complications, local antimicrobial resistance and surveillance data, recent antibiotic use and microbiological results.</p> <p>* Stop antibiotics after 5 days unless microbiological results suggest a longer course is needed or the person is not clinically stable.</p> <p><i>For detailed information click on the visual summary. See also the NICE guideline on pneumonia.</i></p>	First choice (moderate severity in adults): amoxicillin	500mg TDS (higher doses can be used, see BNF)	-	5 days*	
		AND (if atypical pathogens suspected) clarithromycin OR	500mg BD	-		
		erythromycin (in pregnancy)	500mg QDS	-		
		Alternative first choice (moderate severity in adults): doxycycline OR	200mg on day 1, then 100mg OD	-	5 days*	
		clarithromycin	500mg BD	-		
		First choice (high severity in adults or severe in children): co-amoxiclav	500/125mg TDS		5 days*	
		AND (if atypical pathogens suspected) clarithromycin OR	500mg BD			
		erythromycin (in pregnancy)	500mg QDS			
		Alternative first choice (high severity in adults): levofloxacin* (consider safety issues)	Refer to microbiologist	-		
		IV antibiotics - Refer to/Contact Community IV Team – 01253 951223				
Hospital-acquired pneumonia	If symptoms or signs of pneumonia start within 48 hours of hospital discharge, refer to microbiologist.					


Infection	Key points	Medicine	Doses		Length	Visual summary		
			Adult	Child				
▼ Urinary tract infections								
Lower urinary tract infection NICE Public Health England Last updated: Oct 2018	Advise paracetamol or ibuprofen for pain. Non-pregnant women: back up antibiotic (to use if no improvement in 48 hours or symptoms worsen at any time) or immediate antibiotic. Pregnant women, men, children or young people: immediate antibiotic. When considering antibiotics, take account of severity of symptoms, risk of complications, previous urine culture and susceptibility results, previous antibiotic use which may have led to resistant bacteria and local antimicrobial resistance data. If people have symptoms of pyelonephritis (such as fever) or a complicated UTI, see acute pyelonephritis (upper urinary tract infection) for antibiotic choices. <i>For detailed information click on the visual summary. See also the NICE guideline on urinary tract infection in under 16s: diagnosis and management and the Public Health England urinary tract infection: diagnostic tools for primary care.</i>	Non-pregnant women first choice: nitrofurantoin (if eGFR ≥ 45 ml/minute) OR trimethoprim (if low risk of resistance)	100mg m/r BD (or if unavailable 50mg QDS) 200mg BD	-	3 days			
		Non-pregnant women second choice: nitrofurantoin (if eGFR ≥ 45 ml/minute) OR pivmecillinam (a penicillin) OR	100mg m/r BD (or if unavailable 50mg QDS) 400mg initial dose, then 200mg TDS	-	3 days			
		fosfomycin	3g single dose sachet	-	single dose			
		Pregnant women first choice: nitrofurantoin (avoid at term) – if eGFR ≥ 45 ml/minute	100mg m/r BD (or if unavailable 50mg QDS)	-	7 days			
		Pregnant women second choice: amoxicillin (only if culture results available and susceptible) OR cefalexin	500mg TDS 500mg BD	-	7 days			
		Treatment of asymptomatic bacteriuria in pregnant women: choose from nitrofurantoin (avoid at term), amoxicillin or cefalexin based on recent culture and susceptibility results						
		Men first choice: trimethoprim OR nitrofurantoin (if eGFR ≥ 45 ml/minute)	200mg BD 100mg m/r BD (or if unavailable 50mg QDS)	-	7 days			






Summary of antimicrobial prescribing guidance – managing common infections (October 2020)


Review October 2021


Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
		<p>Men second choice: consider alternative diagnoses basing antibiotic choice on recent culture and susceptibility results</p> <p>Children and young people (3 months and over) first choice: trimethoprim (if low risk of resistance) OR nitrofurantoin (if eGFR ≥45 ml/minute)</p> <p>Children and young people (3 months and over) second choice: nitrofurantoin (if eGFR ≥45 ml/minute and not used as first choice) OR amoxicillin (only if culture results available and susceptible) OR cefalexin</p>	-	-	-	
<p>Acute prostatitis</p> <p>NICE</p> <p>Public Health England</p>	<p>Advise paracetamol (+/- low-dose weak opioid) for pain, or ibuprofen if preferred and suitable. Offer antibiotic.</p> <p>Review antibiotic treatment after 14 days and either stop antibiotics or continue for a further 14 days if needed (based on assessment of history, symptoms, clinical examination, urine and blood tests).</p> <p><i>For detailed information click on the visual summary.</i></p>	<p>First choice (guided by susceptibilities when available): ciprofloxacin* (consider safety advice on page 2) OR ofloxacin (consider safety issues on page 2) OR trimethoprim (if fluoroquinolone not appropriate; seek specialist advice)</p>	500mg BD	-	14 days then review	

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Last updated: Oct 2018		Second choice (after discussion with specialist): levofloxacin* (consider safety issues) OR	Refer to microbiologist	-	14 days then review	
		co-trimoxazole	960mg BD	-		
		IV antibiotics- Refer to/Contact Community IV Team – 01253 951223				

Infection	Key points	Medicine	Doses		Length	Visual summary		
			Adult	Child				
Acute pyelonephritis (upper urinary tract) NICE Public Health England Last updated: Oct 2018	Advise paracetamol (+/- low-dose weak opioid) for pain for people over 12. Offer an antibiotic. When prescribing antibiotics, take account of severity of symptoms, risk of complications, previous urine culture and susceptibility results, previous antibiotic use which may have led to resistant bacteria and local antimicrobial resistance data. Avoid antibiotics that don't achieve adequate levels in renal tissue, such as nitrofurantoin. <i>For detailed information click on the visual summary. See also the NICE guideline on urinary tract infection in under 16s: diagnosis and management and the Public Health England urinary tract infection: diagnostic tools for primary care.</i>	Non-pregnant women and men first choice: cefalexin OR	500mg BD or TDS (up to 1g to 1.5g TDS or QDS for severe infections)	-	7 to 10 days			
		co-amoxiclav (only if culture results available and susceptible) OR	500/125mg TDS	-	7 to 10 days			
		trimethoprim (only if culture results available and susceptible) OR	200mg BD	-	14 days			
		ciprofloxacin (consider safety issues) * on page 2	500mg BD	-	7 days			
		Non-pregnant women and men IV antibiotics - Contact specialist for advice						
		Pregnant women first choice: cefalexin	500mg BD or TDS (up to 1g to 1.5g TDS or QDS for severe infections)	-	7 to 10 days			
		Pregnant women second choice or IV antibiotics - Contact specialist for advice						
		Children and young people (3 months and over) first choice: cefalexin OR	-		-			
		co-amoxiclav (only if culture results available and susceptible)	-					
		Children and young people (3 months and over) IV antibiotics - Contact specialist for advice						







Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Recurrent urinary tract infection NICE Public Health England Last updated Oct 2018	<p>First advise about behavioural and personal hygiene measures, and self-care (with D-mannose or cranberry products) to reduce the risk of UTI. Follow LSCMMG Pathway</p> <p>For postmenopausal women, if no improvement, consider vaginal oestrogen (review within 12 months).</p> <p>For non-pregnant women, if no improvement, consider single-dose antibiotic prophylaxis for exposure to a trigger (review within 6 months).</p> <p>For non-pregnant women (if no improvement or no identifiable trigger) or with specialist advice for pregnant women, men, children or young people, consider a trial of daily antibiotic prophylaxis (review within 6 months).</p> <p>For detailed information click on the visual summary. See also the NICE guideline on urinary tract infection in under 16s: diagnosis and management and the Public Health England urinary tract infection: diagnostic tools for primary care.</p> <p><i>Please note - NICE recommendations adapted on local decisions</i></p> <p>Review with mid-stream urine</p>	<p>First choice antibiotic prophylaxis: trimethoprim (avoid in pregnancy) OR</p>	200mg single dose when exposed to a trigger or 100mg at night		- Review if choice is still appropriate after 3 months-switch to alternative agent	
		nitrofurantoin (avoid at term) - if eGFR ≥ 45 ml/minute	100mg single dose when exposed to a trigger or 50 to 100mg at night		Review if choice is still appropriate after 3 months-switch to alternative agent	
		<p>Second choice antibiotic prophylaxis: amoxicillin OR</p>	500mg single dose when exposed to a trigger or 250mg at night		Review if choice is still appropriate after 3 months-switch to alternative agent	
		cefalexin	500mg single dose when exposed to a trigger or 125mg at night		Review if choice is still appropriate after 3 months-switch to alternative agent	

Infection	Key points	Medicine	Doses		Length	Visual summary		
			Adult	Child				
Catheter-associated urinary tract infection NICE Public Health England Last updated: Nov 2018	<p>Antibiotic treatment is not routinely needed for asymptomatic bacteriuria in people with a urinary catheter. Consider removing or, if not possible, changing the catheter if it has been in place for more than 7 days. But do not delay antibiotic treatment.</p> <p>Advise paracetamol for pain.</p> <p>Advise drinking enough fluids to avoid dehydration.</p> <p>Offer an antibiotic for a symptomatic infection.</p> <p>When prescribing antibiotics, take account of severity of symptoms, risk of complications, previous urine culture and susceptibility results, previous antibiotic use which may have led to resistant bacteria and local antimicrobial resistance data.</p> <p>Do not routinely offer antibiotic prophylaxis to people with a short-term or long-term catheter.</p> <p><i>For detailed information click on the visual summary. See also the Public Health England urinary tract infection: diagnostic tools for primary care.</i></p> <p><i>Please note - NICE recommendations adapted on local decisions</i></p>	Non-pregnant women and men first choice if no upper UTI symptoms: nitrofurantoin (if eGFR ≥45 ml/minute) OR	100mg m/r BD (or if unavailable 50mg QDS)	-	7 days			
		trimethoprim (if low risk of resistance) OR	200mg BD	-				
		amoxicillin (only if culture results available and susceptible)	500mg TDS	-				
		Non-pregnant women and men second choice if no upper UTI symptoms: pivmecillinam (a penicillin)	400mg initial dose, then 200mg TDS	-	7 days			
		Non-pregnant women and men first choice if upper UTI symptoms: cefalexin OR	500mg BD or TDS (up to 1g to 1.5g TDS or QDS for severe infections)	-	7 to 10 days			
		co-amoxiclav (only if culture results available and susceptible) OR	500/125mg TDS	-				
		trimethoprim (only if culture results available and susceptible) OR	200mg BD	-	14 days			
		Ciprofloxacin* (consider safety advice on page 2))	500mg BD	-	7 days			
		Non-pregnant women and men IV antibiotics – Refer to/Contact Community IV Team – 01253 951223						
		Pregnant women first choice: cefalexin	500mg BD or TDS (up to 1g to 1.5g TDS or QDS for severe infections)	-	7 to 10 days			

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
		Pregnant women second choice or IV antibiotics – Contact specialist for advice				
		Children and young people (3 months and over) first choice: trimethoprim (if low risk of resistance) OR	-			
		amoxicillin (only if culture results available and susceptible) OR	-			
		cefalexin OR	-			
		co-amoxiclav (only if culture results available and susceptible)	-			
		Children and young people (3 months and over) IV antibiotics- Refer to specialist				
▼ Meningitis						
Suspected meningococcal disease Public Health England Last updated: Feb 2019	Transfer all patients to hospital immediately. ^{1D} If time before hospital admission, ^{2D,3A+} if suspected meningococcal septicaemia or non-blanching rash, ^{2D,4D} give IM benzylpenicillin ^{1D,2D,4D} as soon as possible. ^{2D} Do not give IV/IM antibiotics if there is a definite history of anaphylaxis; ^{1D} rash is not a contraindication. ^{1D}	IM benzylpenicillin ^{1D,2D}	Child <1 year: 300mg ^{5D} Child 1 to 9 years: 600mg ^{5D} Adult/child 10+ years: 1.2g ^{5D}	Stat dose; ^{1D} give IM, if vein cannot be accessed ^{1D}		<i>Not available. Access the supporting evidence and rationales on the PHE website</i>
Prevention of secondary case of meningitis Public Health England Last updated: July 2019	Only prescribe following advice from your local health protection specialist- North West Team: ☎ [0344 225 0562] Out of hours: ☎ [0151 434 4819] Blackpool Teaching Hospitals, microbiologist secretary ☎ [01253 957141] Bleep - 774 Out of hours: contact on-call doctor: ☎ [01253 300000] Expert advice is available for managing clusters of meningitis. Please alert the appropriate organisation to any cluster situation. Public Health England, Colindale (tel: 0208 200 4400) <i>Access the supporting evidence and rationales on the PHE website.</i>					







Summary of antimicrobial prescribing guidance – managing common infections (October 2020)

Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
▼ Gastrointestinal tract infections						
Oral candidiasis Public Health England Last updated: Oct 2018	Topical azoles are more effective than topical nystatin. ^{1A+} Oral candidiasis is rare in immunocompetent adults; ^{2D} consider undiagnosed risk factors, including HIV. ^{2D} Use 50mg fluconazole if extensive/severe candidiasis; ^{3D,4D} if HIV or immunocompromised, use 100mg fluconazole. ^{3D,4D}	Miconazole oral gel ^{1A+,4D,5A-}	2.5ml of 24mg/ml QDS (hold in mouth after food) ^{4D}		7 days; continue for 7 days after resolved ^{4D,6D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		If not tolerated: nystatin suspension ^{2D,6D,7A-}	1ml; 100,000units/ml QDS (half in each side) ^{2D,4D,7A-}		7 days; continue for 2 days after resolved ^{4D}	
		fluconazole capsules ^{6D,7A-}	50mg/100mg OD ^{3D,6D,8A-}		7 to 14 days ^{6D,7A-,8A-}	
Infectious diarrhoea Public Health England Last updated: Oct 2018	Refer previously healthy children with acute painful or bloody diarrhoea, to exclude <i>E. coli</i> O157 infection. ^{1D} Antibiotic therapy is not usually indicated unless patient is systemically unwell. ^{2D} If systemically unwell and campylobacter suspected (such as undercooked meat and abdominal pain), ^{3D} consider clarithromycin 250mg to 500mg BD for 5 to 7 days, if treated early (within 3 days). ^{3D,4A+} If giardia is confirmed or suspected – tinidazole 2g single dose is the treatment of choice. ^{5A+} <i>Access the supporting evidence and rationales on the PHE website.</i>					
Clostridium difficile Public Health England Last updated: Oct 2018	Review need for antibiotics, ^{1D,2D} PPIs, ^{3B-} and antiperistaltic agents and discontinue use where possible. ^{2D} Mild cases (<4 episodes of stool/day) may respond without metronidazole; ^{2D} 70% respond to metronidazole in 5 days; 92% respond to metronidazole in 14 days. ^{4B-} If severe (T>38.5, or WCC>15, rising creatinine, or signs/symptoms of severe colitis): ^{2D} treat with oral vancomycin, ^{1D,2D,5A-} review progress closely, ^{1D,2D} and consider hospital referral. ^{2D}	First episode: metronidazole ^{2D,4B-}	400mg TDS ^{1D,2D}		10 to 14 days ^{1D,4B-}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		Severe, type 027 or recurrent: oral vancomycin ^{1D,2D,5A-}	125mg QDS ^{1D,2D,5A-}		10 to 14 days, ^{1D,2D} then taper ^{2D}	
		Recurrent or second line: (Prescribed in secondary care - can continue on specialist advice) fidaxomicin ^{2D,5A-}	200mg BD ^{5A-}	-	10 days ^{5A-}	
Helicobacter pylori Public Health	Always test for <i>H.pylori</i> before giving antibiotics. Do not offer eradication for GORD. ^{3D} Do not use clarithromycin, metronidazole or quinolone if used in the past year for any infection. ^{5A+,6B+,7A+}	Always use PPI ^{2D,3D,5A+,12A+} First line and first relapse and no penicillin allergy PPI PLUS 2 antibiotics	-		7 days ^{2D} MALToma 14 days ^{7A+,16A+}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>


Summary of antimicrobial prescribing guidance – managing common infections (October 2020)

Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
England See PHE quick reference guide for diagnostic advice: PHE <i>H. pylori</i> Last updated: Feb 2019	<p>Penicillin allergy: use PPI PLUS clarithromycin PLUS metronidazole.^{2D} If previous clarithromycin, use PPI PLUS bismuth salt PLUS metronidazole PLUS tetracycline hydrochloride.^{2D,8A-,9D}</p> <p>Relapse and no penicillin allergy use PPI PLUS amoxicillin PLUS clarithromycin or metronidazole (whichever was not used first line)^{2D}</p> <p>Relapse and previous metronidazole and clarithromycin: use PPI PLUS amoxicillin PLUS either tetracycline OR levofloxacin (if tetracycline not tolerated).^{2D,7A+}</p> <p>Relapse and penicillin allergy (no exposure to quinolone): use PPI PLUS metronidazole PLUS levofloxacin.^{2D}</p> <p>Relapse and penicillin allergy (with exposure to quinolone): use PPI PLUS bismuth salt PLUS metronidazole PLUS tetracycline.^{2D}</p> <p>Retest for <i>H. pylori</i>: post DU/GU, or relapse after second-line therapy,^{1A+} using UBT or SAT,^{10A+,11A+} consider referral for endoscopy and culture.^{2D}</p> <p><i>Please note - NICE recommendations adapted on local decisions</i></p>	amoxicillin ^{2D,6B+} PLUS	1000mg BD ^{14A+}			website
		clarithromycin ^{2D,6B+} OR	500mg BD ^{8A-}			
		metronidazole ^{2D,6B+}	400mg BD ^{2D}			
		Penicillin allergy and previous clarithromycin: PPI WITH bismuth subsalicylate PLUS 2 antibiotics	-	-		
		bismuth subsalicylate ^{13A+} PLUS	525mg QDS ^{15D}			
		metronidazole ^{2D} PLUS	400mg BD ^{2D}			
		tetracycline ^{2D}	500mg QDS ^{15D}			
		Relapse and previous metronidazole and clarithromycin: PPI PLUS 2 antibiotics	-	-		
		amoxicillin ^{2D,7A+} PLUS	1000mg BD ^{14A+}			
		tetracycline ^{2D,7A+} OR	500mg QDS ^{15D}			
		levofloxacin (if tetracycline cannot be used) ^{2D,7A+} *Consider safety issues	250mg BD ^{7A+}			
		Third line- Seek specialist advice				
Acute diverticulitis	<p>Acute diverticulitis and systemically well: Consider no antibiotics, offer simple analgesia (for example paracetamol), advise to re-present if symptoms persist or worsen.</p> <p>Acute diverticulitis and systemically unwell, immunosuppressed or significant</p>	First-choice (uncomplicated acute diverticulitis): co-amoxiclav	500/125mg TDS	-	5 days**	

Summary of antimicrobial prescribing guidance – managing common infections (October 2020)

Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
NICE Last updated: Nov 2019	comorbidity: offer an antibiotic. Give oral antibiotics if person not referred to hospital for suspected complicated acute diverticulitis. Give IV antibiotics if admitted to hospital with suspected or confirmed complicated acute diverticulitis (including diverticular abscess). If CT-confirmed uncomplicated acute diverticulitis, review the need for antibiotics. ** A longer course may be needed based on clinical assessment. <i>Please note - NICE recommendations adapted on local decisions</i>	Penicillin allergy or co-amoxiclav unsuitable: cefalexin (caution in penicillin allergy) AND metronidazole OR	cefalexin: 500mg BD or TDS (up to 1g to 1.5g TDS or QDS for severe infections) metronidazole: 400mg TDS	-		
		trimethoprim AND metronidazole OR	trimethoprim: 200mg BD metronidazole: 400mg TDS	-		
		ciprofloxacin (only if switching from IV ciprofloxacin with specialist advice; consider safety issues) AND metronidazole	ciprofloxacin: 500mg BD metronidazole: 400mg TDS			
		For IV antibiotics in complicated acute diverticulitis (including diverticular abscess) – Refer to/Contact Community IV Team – 01253 951223				
Threadworm Public Health England Last updated: Nov 2017	Treat all household contacts at the same time. ^{1D} . For patients over 2 years old, refer to pharmacy. Advise hygiene measures for 2 weeks ^{1D} (hand hygiene; ^{2D} pants at night; morning shower, including perianal area). ^{1D,2D} Wash sleepwear, bed linen, and dust and vacuum. ^{1D} Child <6 months , add perianal wet wiping or washes 3 hourly. ^{1D} <i>Please note - NICE recommendations adapted on local decisions</i>	Child over 6 months and under 2 years: mebendazole ^{1D,3B-}	100mg stat ^{3B-}		1 dose; ^{3B-} repeat in 2 weeks if persistent ^{3B-}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		Child <6 months or pregnant (at least in first trimester): only hygiene measure for 6 weeks ^{1D}	-	-	-	

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
<p>▼ Genital tract infections</p> <p>For referrals to the GUM clinic between 9am to 5pm ☎ [01253 956850] or ☎ [01253 956931] Out of hours: contact on-call doctor: ☎ [01253 300000]</p>						
<p>STI screening Public Health England Last updated: Nov 2017</p>	<p>People with risk factors should be screened for chlamydia, gonorrhoea, HIV and syphilis.^{1D} Refer individual and partners to GUM.^{1D} Risk factors: <25 years; no condom use; recent/frequent change of partner; symptomatic or infected partner; area of high HIV.^{2B-} Access the supporting evidence and rationales on the PHE website.</p>					

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Chlamydia trachomatis/ urethritis Public Health England Last updated: July 2019	<p>Opportunistically screen all sexually active patients aged 15 to 24 years for chlamydia annually and on change of sexual partner.^{1B-} If positive, treat index case, refer to GUM and initiate partner notification, testing and treatment.^{2D,3A+}</p> <p>As single dose azithromycin has led to increased resistance in GU infections, doxycycline should be used first line for chlamydia and urethritis.^{4A+}</p> <p>Advise patient with chlamydia to abstain from sexual intercourse until doxycycline is completed or for 7 days after treatment with azithromycin (14 days after azithromycin started and until symptoms resolved if urethritis).^{3A+,4A+}</p> <p>If chlamydia, test for reinfection at 3 to 6 months following treatment if under 25 years; or consider if over 25 years and high risk of re-infection.^{1B-,3B+, 5B-}</p> <p>Second line, pregnant, breastfeeding, allergy, or intolerance: azithromycin is most effective.^{6A+,7D,8A+,9A+,10D} As lower cure rate in pregnancy, test for cure at least 3 weeks after end of treatment.^{3A+}</p> <p>Consider referring all patients with symptomatic urethritis to GUM as testing should include <i>Mycoplasma genitalium</i> and <i>Gonorrhoea</i>.^{11A-}</p> <p>If <i>M.genitalium</i> is proven, use doxycycline followed by azithromycin using the same dosing regimen and advise to avoid sex for 14 days after start of treatment and until symptoms have resolved.^{11A-,12A+}</p>	First line: doxycycline ^{4A+,11A-,12A+}	100mg BD ^{4A+,11A-,12A+}		7 days ^{4A+,11A-,12A+}	Not available. Access supporting evidence and rationales on the PHE website
		Second line/ pregnant/breastfeeding/ allergy/intolerance: azithromycin ^{4A+,11A-,12A+}	1000mg ^{4A+,11A-,12A+} then 500mg OD ^{4A+,11A-,12A+}		Stat ^{4A+,11A-,12A+} 2 days ^{4A+,11A-,12A+} (total 3 days)	



Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Epididymitis Public Health England Last updated: Nov 2017	Usually due to Gram-negative enteric bacteria in men over 35 years with low risk of STI. ^{1A+,2D} If under 35 years or STI risk, refer to GUM. ^{1A+,2D}	Doxycycline ^{1A+,2D} OR	100mg BD ^{1A+,2D}	-	10 to 14 days ^{1A+,2D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		ofloxacin ^{1A+,2D} OR	200mg BD ^{1A+,2D}		14 days ^{1A+,2D}	
		ciprofloxacin *(Consider safety advice on page 2) ^{1A+,2D}	500mg BD ^{1A+,2D,3A+}		10 days ^{1A+,2D,3A+}	
Vaginal candidiasis Public Health England Last updated: Oct 2018	All topical and oral azoles give over 80% cure. ^{1A+,2A+} Pregnant: avoid oral azoles, the 7 day courses are more effective than shorter ones. ^{1A+,3D,4A+} Recurrent (>4 episodes per year): ^{1A+} 150mg oral fluconazole every 72 hours for 3 doses induction, ^{1A+} followed by 1 dose once a week for 6 months maintenance. ^{1A+}	Clotrimazole ^{1A+,5D} OR	500mg pessary ^{1A+}	-	Stat ^{1A+}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		fenticonazole ^{1A+} OR	600mg pessary ^{1A+}		Stat ^{1A+}	
		clotrimazole ^{1A+} OR	100mg pessary ^{1A+}		6 nights ^{1A+}	
		oral fluconazole ^{1A+,3D}	150mg ^{1A+,3D}		Stat ^{1A+}	
		If recurrent: fluconazole (induction/maintenance) ^{1A+}	150mg every 72 hours THEN 150mg once a week ^{1A+,3D}		3 doses 6 months ^{1A+}	
Bacterial vaginosis Public Health England Last updated: Nov 2017	Oral metronidazole is as effective as topical treatment, ^{1A+} and is cheaper. ^{2D} 7 days results in fewer relapses than 2g stat at 4 weeks. ^{1A+,2D} Pregnant/breastfeeding: avoid 2g dose. ^{3A+,4D} Treating partners does not reduce relapse. ^{5A+}	oral metronidazole ^{1A+,3A+} OR	400mg BD ^{1A+,3A+} OR 2000mg ^{1A+,2D}	-	7 days ^{1A+} OR Stat ^{2D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		metronidazole 0.75% vaginal gel ^{1A+,2D,3A+} OR	5g applicator at night ^{1A+,2D,3A+}		5 nights ^{1A+,2D,3A+}	
		clindamycin 2% cream ^{1A+,2D}	5g applicator at night ^{1A+,2D}		7 nights ^{1A+,2D,3A+}	
Genital herpes Public Health England Last updated: Nov 2017	Advise: saline bathing, ^{1A+} analgesia, ^{1A+} or topical lidocaine for pain, ^{1A+} and discuss transmission. ^{1A+} First episode: treat within 5 days if new lesions or systemic symptoms, ^{1A+,2D} and refer to GUM. ^{2D} Recurrent: self-care if mild, ^{2D} or immediate short course antiviral treatment, ^{1A+,2D} or suppressive therapy if more than 6 episodes per year. ^{1A+,2D}	oral aciclovir ^{1A+,2D,3A+,4A+} OR	400mg TDS ^{1A+,3A+} 800mg TDS (if recurrent) ^{1A+}	-	5 days ^{1A+} 2 days ^{1A+}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		valaciclovir ^{1A+,3A+,4A+} OR	500mg BD ^{1A+}		5 days ^{1A+}	
		famciclovir ^{1A+,4A+}	250mg TD ^{1A+}		5 days ^{1A+}	
			1000mg BD (if recurrent) ^{1A+}		1 day ^{1A+}	

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Gonorrhoea Public Health England Last updated: Feb 2019	Refer to GUM. ^{3B-} Test of cure is essential. ^{2D} Antibiotic resistance is now very high. ^{1D,2D} Use Ciprofloxacin only If susceptibility is known prior to treatment and the isolate is sensitive to ciprofloxacin at all sites of infection ^{1D,2D} For sensitivity to other antibiotics refer to GUM	Refer to GUM Ciprofloxacin ^{2D} (only if known to be sensitive) *consider safety advice on page 2	500mg ^{2D}	-	Stat ^{2D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
Trichomoniasis Public Health England Last updated: Nov 2017	Oral treatment needed as extravaginal infection common. ^{1D} Treat partners, ^{1D} and refer to GUM for other STIs. ^{1D} Pregnant/breastfeeding: avoid 2g single dose metronidazole; ^{2A+,3D} clotrimazole for symptom relief (not cure) if metronidazole declined. ^{2A+,4A-,5D}	metronidazole ^{1A+,2A+,3D,6A+} Pregnancy to treat symptoms: clotrimazole ^{2A+,4A-,5D}	400mg BD ^{1A+,6A+} 2g (more adverse effects) ^{6A+} 100mg pessary at night ^{5D}	-	5 to 7 day ^{1A+} Stat ^{1A+,6A+} 6 nights ^{5D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
Pelvic inflammatory disease Public Health England Last updated: Feb 2019	Refer women and sexual contacts to GUM. ^{1A+} Raised CRP supports diagnosis, absent pus cells in HVS smear good negative predictive value. ^{1A+} Exclude: ectopic pregnancy, appendicitis, endometriosis, UTI, irritable bowel, complicated ovarian cyst, functional pain. Moxifloxacin has greater activity against likely pathogens, but always test for gonorrhoea, chlamydia, and <i>M. genitalium</i> . ^{1A+} <i>If M. genitalium</i> tests positive use moxifloxacin. ^{1A+}	First line therapy: ceftriaxone ^{1A+,3C,4C} PLUS metronidazole ^{1A+,5A+} PLUS doxycycline ^{1A+,5A+} Second line therapy: metronidazole ^{1A+,5A+} PLUS ofloxacin ^{1A+,2A-,5A+} OR moxifloxacin alone ^{1A+} (first line for <i>M. genitalium</i> associated PID)	1000mg IM ^{1A+,3C} 400mg BD ^{1A+} 100mg BD ^{1A+} 400mg BD ^{1A+} 400mg BD ^{1A+,2A-} Refer to GUM	-	Stat ^{1A+,3C} 14 days ^{1A+} 14 days ^{1A+} 14 days ^{1A+} 14 days ^{1A+}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
▼ Skin and soft tissue infections						
<i>Note: Refer to RCGP Skin Infections online training.^{1D} For MRSA, discuss therapy with microbiologist.^{1D}</i>						
Cold sores Public Health England Last updated: Nov 2017	Most resolve after 5 days without treatment. ^{1A-,2A-} Topical antivirals applied prodromal can reduce duration by 12 to 18 hours. ^{1A-,2A-,3A-} If frequent, severe, and predictable triggers: consider oral prophylaxis: ^{4D,5A+} aciclovir 400mg, twice daily, for 5 to 7 days. ^{5A+,6A+} Access supporting evidence and rationales on the PHE website					

Summary of antimicrobial prescribing guidance – managing common infections (October 2020)








Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
PVL-SA Public Health England Last updated: Nov 2017	Panton-Valentine leukocidin (PVL) is a toxin produced by 20.8 to 46% of <i>S. aureus</i> from boils/abscesses. ^{1B+,2B+,3B-} PVL strains are rare in healthy people, but severe. ^{2B+} Suppression therapy should only be started after primary infection has resolved, as ineffective if lesions are still leaking. ^{4D} Risk factors for PVL: recurrent skin infections; ^{2B+} invasive infections; ^{2B+} MSM; ^{3B-} if there is more than one case in a home or close community ^{2B+,3B-} (school children; ^{3B-} military personnel; ^{3B-} nursing home residents; ^{3B-} household contacts). ^{3B-} Access the supporting evidence and rationales on the PHE website .					
Eczema Public Health England Last updated: Nov 2017	No visible signs of infection: antibiotic use (alone or with steroids) ^{1A+} encourages resistance and does not improve healing. ^{1A+} With visible signs of infection: use oral flucloxacillin ^{2D} or clarithromycin, ^{2D} or topical treatment (as in impetigo). ^{2D} Access the supporting evidence and rationales on the PHE website .					

Infection	Key points	Medicine	Doses		Length	Visual summary	
			Adult	Child			
Impetigo NICE Public Health England Last updated: Feb 2020	Localised non-bullous impetigo: Hydrogen peroxide 1% cream (other topical antiseptics are available but no evidence for impetigo). If hydrogen peroxide unsuitable or ineffective, short-course topical antibiotic. Widespread non-bullous impetigo: Short-course topical or oral antibiotic. Take account of person's preferences, practicalities of administration, previous use of topical antibiotics because antimicrobial resistance can develop rapidly with extended or repeated use, and local antimicrobial resistance data. Bullous impetigo, systemically unwell, or high risk of complications: Short-course oral antibiotic. Do not offer combination treatment with a topical and oral antibiotic to treat impetigo. *5 days is appropriate for most, can be increased to 7 days based on clinical judgement. <i>For detailed information click on the visual summary.</i>	Topical antiseptic:				5 days*	
		hydrogen peroxide 1%	BD or TDS				
		Topical antibiotic:				5 days*	
		First choice: fusidic acid 2%	TDS				
		Fusidic acid resistance suspected or confirmed: mupirocin 2%	TDS				
		Oral antibiotic:				5 days*	
		First choice: flucloxacillin	500mg QDS				
		Penicillin allergy or flucloxacillin unsuitable: clarithromycin OR erythromycin (in pregnancy)	250mg BD 250 to 500mg QDS				
If MRSA suspected or confirmed – consult local microbiologist							
Leg ulcer infection NICE Public Health England	Manage any underlying conditions to promote ulcer healing. Only offer an antibiotic when there are symptoms or signs of infection (such as redness or swelling spreading beyond the ulcer, localised warmth, increased pain or fever). Few leg ulcers are clinically infected but most are colonised by bacteria. When prescribing antibiotics, take account of	First-choice:					
		flucloxacillin	500mg to 1g QDS	-	7 days		
		Penicillin allergy or if flucloxacillin unsuitable:					
		doxycycline OR	200mg on day 1, then 100mg OD (can be increased to 200mg daily)	-	7 days		
		clarithromycin OR erythromycin (in pregnancy)	500mg BD 500mg QDS				





Summary of antimicrobial prescribing guidance – managing common infections (October 2020)

Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary		
			Adult	Child				
Last updated: Feb 2020	severity, risk of complications and previous antibiotic use. <i>For detailed information click on the visual summary.</i>	Second choice:						
		co-amoxiclav OR	500/125mg TDS	-	7 days			
		co-trimoxazole (in penicillin allergy)	960mg BD					
For antibiotic choices if severely unwell or MRSA suspected or confirmed, click on the visual summary								
Cellulitis and erysipelas NICE Public Health England Last updated: Sept 2019	Exclude other causes of skin redness (inflammatory reactions or non-infectious causes). Consider marking extent of infection with a single-use surgical marker pen. Offer an antibiotic. Take account of severity, site of infection, risk of uncommon pathogens, any microbiological results and MRSA status. Infection around eyes or nose is more concerning because of serious intracranial complications. *A longer course (up to 14 days in total) may be needed but skin takes time to return to normal, and full resolution at 5 to 7 days is not expected. Do not routinely offer antibiotics to prevent recurrent cellulitis or erysipelas. <i>For detailed information click on the visual summary.</i>	First choice:						
		flucloxacillin	500mg to 1g QDS		5 to 7 days*			
		Penicillin allergy or if flucloxacillin unsuitable:						
		clarithromycin OR	500mg BD		5 to 7 days*			
		erythromycin (in pregnancy) OR	500mg QDS					
		doxycycline (adults only) OR	200mg on day 1, then 100mg OD	-				
		co-amoxiclav (children only: not in penicillin allergy)	-					
		If infection near eyes or nose:						
		co-amoxiclav	500/125mg TDS		7 days*			
		If infection near eyes or nose (penicillin allergy):						
clarithromycin AND	500mg BD		7 days*					
metronidazole (only add in children if anaerobes suspected)	400mg TDS							
For alternative choice antibiotics for severe infection, suspected or confirmed MRSA infection and IV antibiotics contact local microbiologist								
Diabetic foot infection NICE	In diabetes, all foot wounds are likely to be colonised with bacteria. Diabetic foot infection has at least 2 of: local swelling or induration; erythema; local tenderness or pain; local warmth; purulent discharge. Severity is classified as:	Mild infection: first choice						
		flucloxacillin	500mg to 1g QDS	-	7 days*			
		Mild infection (penicillin allergy):						
		clarithromycin OR	500mg BD	-	7 days*			
erythromycin (in pregnancy) OR	500mg QDS							











Summary of antimicrobial prescribing guidance – managing common infections (October 2020)






Review October 2021





Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Public Health England Last updated: Oct 2019	<p>Mild: local infection with 0.5 to less than 2cm erythema</p> <p>Moderate: local infection with more than 2cm erythema or involving deeper structures (such as abscess, osteomyelitis, septic arthritis or fasciitis)</p> <p>Severe: local infection with signs of a systemic inflammatory response.</p> <p>Start antibiotic treatment as soon as possible.</p> <p>Take samples for microbiological testing before, or as close as possible to, the start of treatment</p> <p>When choosing an antibiotic, take account of severity, risk of complications, previous microbiological results and antibiotic use, and patient preference.</p> <p>*A longer course (up to a further 7 days) may be needed based on clinical assessment. However, skin does take time to return to normal, and full resolution at 7 days is not expected.</p> <p>Do not offer antibiotics to prevent diabetic foot infection.</p> <p><i>For detailed information click on the visual summary.</i></p>	doxycycline	200mg on day 1, then 100mg OD (can be increased to 200mg daily)			
<p>For antibiotic choices for moderate or severe infection, infections where <i>Pseudomonas aeruginosa</i> or MRSA is suspected or confirmed, and IV antibiotics contact local microbiologist</p>						
<p>Tick bites (Lyme disease)</p> <p>Public Health England</p> <p>Last updated: Feb 2020</p>	<p>Treatment: Treat erythema migrans empirically; serology is often negative early in infection.^{1D}</p> <p>For other suspected Lyme disease such as neuroborreliosis (CN palsy, radiculopathy) seek advice.^{1D}</p>	<p>Treatment: doxycycline^{1D}</p> <hr/> <p>Alternative: amoxicillin^{1D}</p>	100mg BD ^{1D}		21 days ^{1D}	Not available. Access supporting evidence and rationales on the PHE website
			1,000mg TDS ^{1D}			
Acne	<p>Mild (open and closed comedones)^{1D} or moderate (inflammatory lesions):^{1D}</p> <p>First line: self-care^{1D} (wash with mild soap; do not scrub; avoid make-up).^{1D}</p>	Second line: topical retinoid ^{1D,2D,3A+} OR	Thinly OD ^{3A+}		6 to 8 weeks ^{1D}	Not available. Access supporting evidence and rationales on
		benzoyl peroxide ^{1A-,2D,3A+,4A-}	5% cream OD-BD ^{3A+}		6 to 8 weeks ^{1D}	




Summary of antimicrobial prescribing guidance – managing common infections (October 2020)

Review October 2021

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Public Health England Last updated: Nov 2017	Second line: topical retinoid or benzoyl peroxide. ^{2D} Third-line: add topical antibiotic, ^{1D,3A+} or consider addition of oral antibiotic. ^{1D} Severe (nodules and cysts): ^{1D} add oral antibiotic (for 3 months max) ^{1D,3A+} and refer. ^{1D,2D}	Third-line: topical clindamycin ^{3A+}	1% cream, thinly BD ^{3A+}		12 weeks ^{1A-,2D}	the PHE website
		If treatment failure/severe: oral tetracycline ^{1A-,3A+} OR	500mg BD ^{3A+}		6 to 12 weeks ^{3A+}	
		oral doxycycline ^{3A+,4A-}	100mg OD ^{3A+}		6 to 12 weeks ^{3A+}	
Scabies Public Health England Last updated: Oct 2018	First choice permethrin: Treat whole body from ear/chin downwards, ^{1D,2D} and under nails. ^{1D,2D} If using permethrin and patient is under 2 years, elderly or immunosuppressed, or if treating with malathion: also treat face and scalp. ^{1D,2D} Home/sexual contacts: treat within 24 hours. ^{1D}	permethrin ^{1D,2D,3A+}	5% cream ^{1D,2D}		2 applications, 1 week apart ^{1D}	Not available. Access supporting evidence and rationales on the PHE website
		Permethrin allergy: malathion ^{1D}	0.5% aqueous liquid ^{1D}			
Bites Public Health England Last updated: July 2019	Human: thorough irrigation is important. ^{1A+,2D} Antibiotic prophylaxis is advised. ^{1A+,2D,3D} Assess risk of tetanus, rabies, ^{1A+} HIV, and hepatitis B and C. ^{3D} Cat: always give prophylaxis. ^{1A+,3D} Dog: give prophylaxis if: puncture wound; ^{1A+,3D} bite to hand, foot, face, joint, tendon, or ligament; ^{1A+} immunocompromised; cirrhotic; asplenic; or presence of prosthetic valve/joint. ^{2D,4A+} Penicillin allergy: Review all at 24 and 48 hours, ^{3D} as not all pathogens are covered. ^{2D,3}	Prophylaxis/treatment all: co-amoxiclav ^{2D,3D}	375mg to 625mg TDS ^{3D}		7 days ^{3D}	Not available. Access supporting evidence and rationales on the PHE website
		Human + penicillin allergy: metronidazole ^{3D,4A+} AND	400mg TDS ^{2D}		7 days ^{3D}	
		clarithromycin ^{3D,4A+}	250mg to 500mg BD ^{2D}			
		Animal + penicillin allergy: metronidazole ^{3D,4A+} AND	400mg TDS ^{2D}		7 days ^{3D}	
		doxycycline ^{3D}	100mg BD ^{2D}			

Infection	Key points	Medicine	Doses		Length	Visual summary	
			Adult	Child			
Mastitis Public Health England Last updated: Nov 2017	<p><i>S. aureus</i> is the most common infecting pathogen.^{1D} Suspect if woman has: a painful breast;^{2D} fever and/or general malaise;^{2D} a tender, red breast.^{2D}</p> <p>Breastfeeding: oral antibiotics are appropriate, where indicated.^{2D,3A+} Women should continue feeding,^{1D,2D} including from the affected breast.^{2D}</p>	flucloxacillin ^{2D} Penicillin allergy: erythromycin ^{2D} OR clarithromycin ^{2D}	500mg QDS ^{2D}	250mg to 500mg QDS ^{2D}	-	10 to 14 days ^{2D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
Dermatophyte infection: skin Public Health England Last updated: Feb 2019	<p>Most cases: use terbinafine as fungicidal, treatment time shorter and more effective than with fungistatic imidazoles or undecenoates.^{1D,2A+} If candida possible, use imidazole.^{4D}</p> <p>If intractable, or scalp: send skin scrapings,^{1D} and if infection confirmed: use oral terbinafine^{1D,3A+,4D} or itraconazole.^{2A+,3A+,5D}</p> <p>Scalp: oral therapy,^{6D} and discuss with specialist.^{1D}</p>	topical terbinafine ^{3A+,4D} OR topical imidazole ^{2A+,3A+} Alternative in athlete's foot: topical undecenoates ^{2A+} (such as Mycota®) ^{2A+}	1% OD to BD ^{2A+}	1% OD to BD ^{2A+}	  	1 to 4 weeks ^{3A+} 4 to 6 weeks ^{2A+,3A+}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
Dermatophyte infection: nail Public Health England Last updated: Oct 2018	<p>Take nail clippings;^{1D} start therapy only if infection is confirmed.^{1D} Oral terbinafine is more effective than oral azole.^{1D,2A+,3A+,4D} Liver reactions 0.1 to 1% with oral antifungals.^{3A+} If candida or non-dermatophyte infection is confirmed, use oral itraconazole.^{1D,3A+,4D} Topical nail lacquer is not as effective.^{1D,5A+,6D}</p> <p>To prevent recurrence: apply weekly 1% topical antifungal cream to entire toe area.^{6D}</p> <p>Children: seek specialist advice.^{4D}</p>	First line: terbinafine ^{1D,2A+,3A+,4D,6D} Second line: itraconazole ^{1D,3A+,4D,6D}	250mg OD ^{1D,2A+,6D}	200mg BD ^{1D,4D}	 	Fingers: 6 weeks ^{1D,6D} Toes: 12 weeks ^{1D,6D} 1 week a month ^{1D} Fingers: 2 courses ^{1D} Toes: 3 courses ^{1D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
Stop treatment when continual, new, healthy, proximal nail growth. ^{6D}							

Infection	Key points	Medicine	Doses		Length	Visual summary	
			Adult	Child			
Varicella zoster/ chickenpox Herpes zoster/ shingles Public Health England Last updated: Oct 2018	Pregnant/immunocompromised/ neonate: seek urgent specialist advice. ^{1D} Chickenpox: consider aciclovir ^{2A+,3A+,4D} if: onset of rash <24 hours, ^{3A+} and 1 of the following: >14 years of age; ^{4D} severe pain; ^{4D} dense/oral rash; ^{4D,5B+} taking steroids; ^{4D} smoker. ^{4D,5B+} Give paracetamol for pain relief. ^{6C} Shingles: treat if >50 years ^{7A+,8D} (PHN rare if <50 years) ^{9B+} and within 72 hours of rash, ^{10A+} or if 1 of the following: active ophthalmic; ^{11D} Ramsey Hunt; ^{4D} eczema; ^{4D} non-truncal involvement; ^{8D} moderate or severe pain; ^{8D} moderate or severe rash. ^{5B+,8D} Shingles treatment if not within 72 hours: consider starting antiviral drug up to 1 week after rash onset, ^{12B+} if high risk of severe shingles ^{12B+} or continued vesicle formation; ^{4D} older age; ^{7A+,8D,12B+} immunocompromised; ^{4D} or severe pain. ^{7D,11B+}	First line for chicken pox and shingles: aciclovir ^{3A+,7A+,10A+,13B+,14A-,15A+}	800mg 5 times daily ^{16A-}		7 days ^{14A-,16A-}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>	
		Second line for shingles if poor compliance: not for children: famciclovir ^{8D,14A-,16A-} OR valaciclovir ^{8D,10A+,14A-}	250mg to 500mg TDS ^{15A+} OR 750mg BD ^{15A+}	-			
			1g TDS ^{14A-}				
▼ Eye infections							
Conjunctivitis Public Health England Last updated: July 2019	First line: bath/clean eyelids with cotton wool dipped in sterile saline or boiled (cooled) water, to remove crusting. ^{1D} Treat only if severe, ^{2A+} as most cases are viral ^{3D} or self-limiting. ^{2A+} Bacterial conjunctivitis: usually unilateral and also self-limiting. ^{2A+,3D} It is characterised by red eye with mucopurulent, not watery discharge. ^{3D} 65% and 74% resolve on placebo by days 5 and 7. ^{4A-,5A+} Third line: fusidic acid as it has less Gram-negative activity. ^{6A-,7D}	Second line: chloramphenicol ^{1D,2A+,4A-,5A+} 0.5% eye drop ^{1D,2A+} OR 1% ointment ^{1D,5A+}	Eye drops: 2 hourly for 2 days, ^{1D,2A+} then reduce frequency ^{1D} to 3 to 4 times daily. ^{1D} Eye ointment: 3 to 4 times daily or once daily at night if using antibiotic eye drops during the day. ^{1D}		48 hours after resolution ^{2A+,7D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>	
		Third line: fusidic acid 1% gel ^{2A+,5A+,6A-}	BD ^{1D,7D}				

Infection	Key points	Medicine	Doses		Length	Visual summary
			Adult	Child		
Blepharitis Public Health England Last updated: Nov 2017	First line: lid hygiene ^{1D,2A+} for symptom control, ^{1D} including: warm compresses; ^{1D,2A+} lid massage and scrubs; ^{1D} gentle washing; ^{1D} avoiding cosmetics. ^{1D} Second line: topical antibiotics if hygiene measures are ineffective after 2 weeks. ^{1D,3A+} Signs of meibomian gland dysfunction,^{3D} or acne rosacea:^{3D} consider oral antibiotics.^{1D}	Second line: topical chloramphenicol ^{1D,2A+,3A-}	1% ointment BD ^{2A+,3D}		6-week trial ^{3D}	<i>Not available. Access supporting evidence and rationales on the PHE website</i>
		Third line: oral oxytetracycline ^{1D,3D} OR oral doxycycline ^{1D,2A+,3D}	500mg BD ^{3D} 250mg BD ^{3D}		4 weeks (initial) ^{3D} 8 weeks (maint) ^{3D}	
			100mg OD ^{3D} 50mg OD ^{3D}		4 weeks (initial) ^{3D} 8 weeks (maint) ^{3D}	
▼ Suspected dental infections in primary care (outside dental settings)						
Derived from the Scottish Dental Clinical Effectiveness Programme (SDCEP) 2013 Guidelines. This guidance is not designed to be a definitive guide to oral conditions, as GPs should not be involved in dental treatment. Patients presenting to non-dental primary care services with dental problems should be directed to their regular dentist, or if this is not possible, to the NHS 111 service (in England), who will be able to provide details of how to access emergency dental care.						
<i>Note: Antibiotics do not cure toothache.^{1D} First-line treatment is with paracetamol^{1D} and/or ibuprofen;^{1D} codeine is not effective for toothache.^{1D}</i>						
▼ Abbreviations						
BD, twice a day; eGFR, estimated glomerular filtration rate; IM, intramuscular; IV, intravenous; MALToma, mucosa-associated lymphoid tissue lymphoma; m/r, modified release; MRSA, methicillin-resistant <i>Staphylococcus aureus</i> ; MSM, men who have sex with men; stat, given immediately; OD, once daily; TDS, 3 times a day; QDS, 4 times a day.						